Malaria: Ghana (Q2/2024)

On track | Not applicable | Improve
Progress | No data | Decline
Not on track

Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	⬇0.090	94	92	63	0.94	⬇84.76
Ahafo		89	90	⬆64	1.02	⬇89.3
Ashanti	0.020	95	84	55	0.95	⬆92.42
Bono	0.060	93	89	⬆60	1.01	⬆93.68
Bono East	0.150	91	91	⬆62	⬇0.82	⬇83.77
Central	0.050	96	95	71	0.96	⬇87.51
Eastern	0.050	90	98	68	0.96	⬇88.37
Greater Accra	⬆0.080	99	91	⬆73	0.94	⬇86.92
North East		96	⬇85	55	0.76	⬇66.89
Northern		94	⬇80	53	⬇0.72	⬇60.52
Oti		96	99	⬆71	⬆1.38	⬇77.53
Savannah	2.030	93	95	⬆53	⬇0.61	⬇50.7
Upper East		98	96	⬇53	0.95	⬆92.49
Upper West		98	96	⬆78	⬆1.45	⬆89.76
Volta	0.400	94	93	74	0.99	⬆90.05
Western		95	95	65	0.92	⬇82.27
Western North		84	99	⬆63	⬆0.88	⬇82.97
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2