## **■ Malaria: Ghana (Q2/2024)**

On track	
Progress	
Not on track	

Not applicable
No data





## **Country indicators**

pos	alaria test itivity ratio ntinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	100,000 persons	under 5 years old	% of pregnant women who slept under an ITN the previous night	using ITN among	received 3 doses	IRS in the last 12
DHIM	IS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

## **Scorecard**

Region	In	Impact		Coverage				
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN		
Ghana	0.090	94	92	63	0.94	84.76		
Ahafo		89	90	64	1.02	89.3		
Ashanti	0.020	95	84	55	0.95	92.42		
Bono	0.060	93	89	60	1.01	93.68		
Bono East	0.150	91	91	62	0.82	83.77		
Central	0.050	96	95	71	0.96	87.51		
Eastern	0.050	90	98	68	0.96	88.37		
Greater Accra	0.080	99	91	73	0.94	86.92		
North East		96	85	55	0.76	66.89		
Northern		94	80	53	0.72	60.52		
Oti		96	99	71	1.38	77.53		
Savannah	2.030	93	95	53	0.61	50.7		
Upper East		98	96	53	0.95	92.49		
Upper West		98	96	78	1.45	89.76		
Volta	0.400	94	93	74	0.99	90.05		
Western		95	95	65	0.92	82.27		
Western North		84	99	63	0.88	82.97		
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2		