

## Scorecard

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	↓0.090	94	92	63	0.94	↓84.76
Ahafo		89	90	↑64	1.02	↓89.3
Ashanti	0.020	95	84	55	0.95	↑92.42
Bono	0.060	93	89	↑60	1.01	↑93.68
Bono East	0.150	91	91	↑62	↓0.82	↓83.77
Central	0.050	96	95	71	0.96	↓87.51
Eastern	0.050	90	98	68	0.96	↓88.37
Greater Accra	↑0.080	99	91	↑73	0.94	↓86.92
North East		96	↓85	55	0.76	↓66.89
Northern		94	↓80	53	↓0.72	↓60.52
Oti		96	99	↑71	↑1.38	↓77.53
Savannah	2.030	93	95	↑53	↓0.61	↓50.7
Upper East		98	96	↓53	0.95	↑92.49
Upper West		98	96	↑78	↑1.45	↑89.76
Volta	0.400	94	93	74	0.99	↑90.05
Western		95	95	65	0.92	↓82.27
Western North		84	99	↑63	↑0.88	↓82.97
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2