■ Malaria: Ghana (Q3/2024)

On track
Progress
Not on track

Not applicable
No data





Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	100,000 persons	under 5 years old	% of pregnant women who slept under an ITN the previous night	using ITN among those with access		IRS in the last 12
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	In	Impact		Coverage				
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN		
Ghana	♦ 3.030	98	92	65	1 96.14	↑ 90.53		
Ahafo		1 99	90	↑ 75	1 99.75	↑ 93.99		
Ashanti		97	83	59	1 92.29	♦ 89.35		
Bono		1 99	₽ 84	1 74	1 01.35	♦ 92.25		
Bono East	↓ 21.900	94	94	63	1 97.88	1 91.42		
Central	↓ 4.970	99	94	♦ 66	1 04.35	1 89.6		
Eastern	↓ 3.670	1 99	98	₽ 62	1 96.20	♦ 87.95		
Greater Accra	↓ 5.920	99	92	♦ 67	1 99.12	1 97.62		
North East		97	1 99	1 65	1 98.60	1 99.15		
Northern		98	1 86	57	1 94.81	1 78.01		
Oti		99	99	75	1 97.54	1 90.21		
Savannah		97	92	57	1 85.97	1 71.44		
Upper East		99	98	1 68	1 102.84	1 94.55		
Upper West	11.650	100	94	1 90	1 02.69	1 96.79		
Volta	↓ 20.280	1 99	₽ 87	♦ 65	1 98.72	1 90.64		
Western		98	95	65	1 85.74	1 89.73		
Western North		1 96	98	61	1 81.59	1 83.96		
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2		