



Highlighted indicators

Stunting	Newborn MR	Maternal MR
38	24	439

Scorecard

Region	Sexual and Reproductive Health		Maternal Health					Newborn Health				Child Health		Nutrition		Monitoring and Evaluation	
			% deliveries by skilled birth attendant	% of pregnant women starting ANC in first trimester	% of pregnant women who received 3 or more doses of IPT during ANC	Institutional Maternal Mortality Ratio (HMSIS-based) Indicator	% of HIV+ children initiated on ARTs	Postnatal visit within 2 wks of delivery	Proportion of HIV exposed infants tested for HIV within 2 months of age	% of low birth weight infants accessing KMC services	Fresh still birth rate	Under 1 year children fully immunized	Proportion of under five children with pneumonia treated with antibiotics	Proportion of <5 children with diarrhoea treated with ORS and Zinc	US Vitamin A (routine)	% of babies initiated on breast feeding within 1 hour of birth	Data completeness
Malawi	22	97	3	54	51	▲74	5	▲74	35	8	16	▲97	88	19	91	91	▲81
Central East Zone	20	96	4	55	71	▲93	6	69	34	▲9	16	90	85	▲26	93	95	▲84
Central Hospital	0	99	0	▲23	▲225	▲95	0	31		17	0	0	0	0	66	0	0
Central West Zone	20	97	3	55	▲63	▲91	4	72	40	▲6	16	▲91	86	▲46	89	89	▲78
North Zone	19	96	3	56	35	▲131	4	▲75	34	▲7	16	▲91	▲87	▲38	93	85	73
South East Zone	26	96	4	53	▲18	▲56	6	77	▲32	▲7	16	▲90	▲92	▲24	93	98	92
South West Zone	25	96	3	51	▲35	▲73	6	80	41	7	18	90	91	▲28	92	▲90	▲80
Source:	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2 (Maternity&KMC reports)	DHIS2	DHIS2 (EPI Dataset)	DHIS2	DHIS2	DHIS2 (EPI Dataset)	DHIS2	DHIS2 (HMIS 15 Dataset)	DHIS2 (HMIS 15 Dataset)