



Highlighted indicators

Stunting	Newborn MR	Maternal MR
38	24	439

Carte de score

Region	Sexual and Reproductive Health			Maternal Health				Newborn Health				Child Health		Nutrition		Monitoring and Evaluation	
	% of adolescent youths (10-19 years) accessing Youth Friendly Health Services	% of women of reproductive age group receiving FP services	% deliveries by skilled birth attendant	% of pregnant women starting ANC in first trimester	% of pregnant women who received 3 or more doses of IPT during ANC	Institutional Maternal Mortality Ratio (HMIS-based) Indicator	% of HIV+ children initiated on ARTs	Postnatal visit within 2 wks of delivery	Proportion of HIV exposed infants tested for HIV within 2 months of age	% of low birth weight infants accessing KMC services	Fresh still birth rate	Under 1 year children fully immunized	Proportion of under five children with pneumonia treated with antibiotics	Proportion of <5 children with diarrhoea treated with ORS and Zinc	US Vitamin A (routine)	% of babies initiated on breast feeding within 1 hour of birth	Data completeness
Malawi	22	97	3	54	51	74	5	74	35	8	16	90	88	19	91	91	82
Central East Zone	20	96	4	55	71	35	6	69	34	9	16	90	85	26	93	95	84
Central Hospital	0	99	0	23	225	95	0	31		12	0	0	0	0	65	0	0
Central West Zone	20	97	3	55	63	81	4	72	40	8	16	91	86	46	89	89	78
North Zone	19	96	3	56	35	121	4	75	34	7	16	91	87	38	93	85	73
South East Zone	26	96	4	53	18	56	6	77	32	7	16	85	92	24	93	88	92
South West Zone	25	96	3	51	26	73	6	80	41	7	18	90	91	28	92	85	80
Source:	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2	DHIS2 (Mortality&KMC reports)	DHIS2	DHIS2 (EPI Dataset)	DHIS2	DHIS2	DHIS2 (EPI Dataset)	DHIS2	DHIS2 (HMIS 15 Dataset)	DHIS2 (HMIS 15 Dataset)