



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of childrens due for measles 2 given ITN
Ghana	0.0	91%	100%	49%	89%	90%
Ahafo						
Ashanti	0.0	84%	100%	39%	91%	96%
Bono	0.0	95%	100%	60%	100%	100%
Bono East						
Central	0.0	95%	100%	52%	86%	84%
Eastern	0.0	91%	100%	48%	99%	96%
Greater Accra	0.0	88%	100%	48%	86%	86%
North East						
Northern	0.0	83%	100%	46%	68%	70%
Oti						
Savannah						
Upper East	0.0	97%	100%	56%	89%	87%
Upper West	0.0	97%	100%	61%	72%	70%
Volta	0.0	91%	100%	52%	98%	98%
Western	0.0	91%	100%	51%	99%	96%
Western North						
Source:	DHIMS 2	DHIMS 2	DHMIS	DHIMS 2	DHIMS 2	DHIMS 2