



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
0%	20.4%	4	52%	50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.0	92%	100%	47%	87%	87%
Ahafo						
Ashanti	0.0	84%	100%	36%	90%	94%
Bono	0.0	97%	100%	56%	98%	96%
Bono East						
Central	0.0	93%	100%	54%	85%	81%
Eastern	0.0	93%	100%	48%	98%	98%
Greater Accra	0.0	86%	100%	47%	↓71%	↓72%
North East						
Northern	0.0	↑88%	100%	↓41%	71%	73%
Oti						
Savannah						
Upper East	0.0	97%	100%	↓47%	↑96%	91%
Upper West	0.0	99%	100%	59%	↑79%	↑82%
Volta	0.0	91%	100%	51%	100%	99%
Western	0.0	94%	100%	50%	98%	93%
Western North						
Source:	DHIMS 2	DHIMS 2	DHMIS	DHIMS 2	DHIMS 2	DHIMS 2