



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
21%	20.4%	4	52%	50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.0	93%	93%	43%	↓70%	↓67%
Ahafo						
Ashanti	0.0	86%	92%	36%	↓64%	↓65%
Bono	0.0	96%	85%	54%	87%	95%
Bono East						
Central	0.0	↑92%	99%	↓42%	73%	↓64%
Eastern	0.0	95%	100%	42%	98%	97%
Greater Accra	0.0	90%	90%	46%	↓25%	↓17%
North East						
Northern	0.0	79%	76%	42%	↓62%	↓57%
Oti						
Savannah						
Upper East	0.0	97%	91%	↓49%	↓75%	↓68%
Upper West	0.0	97%	92%	↓41%	95%	95%
Volta	0.0	↑94%	100%	↓45%	98%	98%
Western	0.0	94%	92%	44%	93%	↓85%
Western North						
Source:	DHIMS 2	DHIMS 2	DHMIS2	DHIMS 2	DHIMS 2	DHIMS 2