■ Malaria: Ghana (Q1/2019)

On track
Progress
Not on track

Not applicable
No data





Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	100,000 persons	under 5 years old	% of pregnant women who slept under an ITN the previous night	using ITN among	received 3 doses	IRS in the last 12
100%	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	In	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN	
Ghana	♦ 0.110	1 93	1 26	1 48	↓ 0.23	66%	
Ahafo	0.000	97	864	50		97%	
Ashanti	0.030	1 87	1 08	1 45	↓ 0.00	₹ 55%	
Bono	♦ 0.450	1 98	1 88	1 64	♦ 0.00	₽ 88%	
Bono East	0.160	97	93	58		78%	
Central	0.140	1 94	1 97	1 45	↓ 0.00	1 82%	
Eastern	↓ 0.060	1 96	1 06	1 48	↓ 0.00	96%	
Greater Accra	♦ 0.140	1 90	1 06	± 51	♦ 0.00	19%	
North East	0.330	64	64	33		69%	
Northern	↓ 0.060	1 79	1 93	1 45	1 2.44	↓ 45%	
Oti	0.000	94	114	41		97%	
Savannah	0.080	84	87	30		86%	
Upper East	0.000	1 98	1 99	1 57	↓ 0.00	67%	
Upper West	♦ 0.120	1 98	1 96	1 40	♦ 0.00	92%	
Volta	♦ 0.140	1 96	1 111	1 56	♦ 0.00	96%	
Western	↓ 0.250	1 94	1 94	1 52	♦ 0.00	87%	
Western North	0.100	95	85	52		92%	
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2	