

Indicateurs nationaux

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
100%	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Carte de score

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	↓0.110	↑93	↑126	↑48	↓0.23	66%
Ahafo	0.000	97	864	50		97%
Ashanti	0.030	↑87	↑108	↑45	↓0.00	↓55%
Bono	↓0.450	↑98	↑88	↑64	↓0.00	↓88%
Bono East	0.160	97	93	58		78%
Central	0.140	↑94	↑97	↑45	↓0.00	↑82%
Eastern	↓0.060	↑96	↑106	↑48	↓0.00	96%
Greater Accra	↑0.140	↑90	↑106	↑51	↓0.00	19%
North East	0.330	64	64	33		69%
Northern	↓0.060	↑79	↑93	↑45	↑2.44	↓45%
Oti	0.000	94	114	41		97%
Savannah	0.080	84	87	30		86%
Upper East	0.000	↑98	↑99	↑57	↓0.00	67%
Upper West	↓0.120	↑98	↑96	↑40	↓0.00	92%
Volta	↓0.140	↑96	↑111	↑56	↓0.00	96%
Western	↓0.250	↑94	↑94	↑52	↓0.00	87%
Western North	0.100	95	85	52		92%
Source:	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2