



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
100%	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.080	95	↓96	50	↑0.82	↑78%
Ahafo	↓0.070	97	↓98	↑55	0.88	↓90%
Ashanti	0.010	91	↓99	45	↑0.85	↑73%
Bono	↑0.140	98	↓72	64	↑0.89	86%
Bono East	0.120	98	94	56	0.73	78%
Central	↑0.000	96	97	↑57	↑0.90	86%
Eastern	↓0.160	96	104	47	↑0.98	96%
Greater Accra	↑0.000	92	101	↑58	↑0.76	↑67%
North East	0.350	↑94	↓41	31	0.79	72%
Northern	0.080	81	↓77	↓39	↓0.46	46%
Oti	↓0.170	95	111	45	0.84	↓85%
Savannah	↓0.150	85	↑102	29	0.89	87%
Upper East	0.000	99	101	↓46	↑0.76	↑77%
Upper West	↓0.230	99	100	↑57	↑0.82	↓84%
Volta	0.100	96	113	↑61	↑0.96	98%
Western	↑0.030	95	↑99	↑58	↑0.95	90%
Western North	0.090	97	87	48	0.97	↓87%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2