



Indicateurs nationaux

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
100%	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Carte de score

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.110	94	97	52	↑0.92	↑88%
Ahafo	0.100	99	96	↑61	↑0.98	↑96%
Ashanti	0.040	89	99	49	↑0.97	↑87%
Bono	↓0.310	101	↑78	↑70	↑0.95	↑96%
Bono East	↓0.170	100	↓80	59	↑0.82	77%
Central	↑0.050	95	95	56	0.95	↑92%
Eastern	↑0.040	97	108	48	0.96	94%
Greater Accra	↓0.290	91	↑116	55	↑0.85	↑84%
North East	↑0.130	↓82	44	↑36	↑0.96	↑85%
Northern	0.110	↓72	↑93	↑46	↑0.68	↑67%
Oti	0.160	96	↓103	46	↑0.98	↑100%
Savannah	↑0.000	↓76	↑125	28	0.94	89%
Upper East	↓0.150	98	↓93	↑64	↑0.97	↑84%
Upper West	↑0.110	97	99	↑64	↑0.94	↑98%
Volta	↓0.360	94	↑125	↓55	0.99	97%
Western	0.040	96	96	54	↑1.00	93%
Western North	0.080	97	84	48	0.99	↑94%
Source:	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2