



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
100%	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.110	95	93	↓43	0.92	↓83%
Ahafo	0.070	98	92	↓53	0.97	95%
Ashanti	0.050	92	↓90	↓41	0.93	↓82%
Bono	↑0.050	98	77	65	0.98	91%
Bono East	↓0.330	99	↑91	58	↑0.93	↑92%
Central	↓0.110	95	91	↓45	0.95	88%
Eastern	↓0.120	98	↓101	45	0.97	90%
Greater Accra	0.320	94	↓93	↓46	0.89	81%
North East	↑0.000	82	↑49	↓17	0.92	↑91%
Northern	↓0.190	76	97	↓31	0.66	↓48%
Oti	0.180	98	102	↓35	0.98	99%
Savannah	0.000	80	↓94	27	0.94	↓82%
Upper East	↑0.000	98	↑100	↓48	0.98	↑93%
Upper West	0.120	98	94	↓49	0.93	96%
Volta	↑0.150	97	↓100	↓44	1.00	96%
Western	0.040	96	98	↓43	0.99	89%
Western North	0.040	97	↑92	47	0.96	97%
Source:	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2