Malaria: Ghana (Q4/2019)

On track
Progress
Not on track
Not applicable
No data
Improve
Decline

Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
100%	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.110	95	93	⬇43	0.92	⬇83%
Ahafo	0.070	98	92	⬇53	0.97	95%
Ashanti	0.050	92	⬇90	⬇41	0.93	⬇82%
Bono	⬆0.050	98	77	65	0.98	91%
Bono East	⬇0.330	99	⬆91	58	⬆0.93	⬆92%
Central	⬇0.110	95	91	⬇45	0.95	88%
Eastern	⬇0.120	98	⬇101	45	0.97	90%
Greater Accra	0.320	94	⬇93	⬇46	0.89	81%
North East	⬆0.000	82	⬆49	⬇17	0.92	⬆91%
Northern	⬇0.190	76	97	⬇31	0.66	⬇48%
Oti	0.180	98	102	⬇35	0.98	99%
Savannah	0.000	80	⬇94	27	0.94	⬇82%
Upper East	⬆0.000	98	⬆100	⬇48	0.98	⬆93%
Upper West	0.120	98	94	⬇49	0.93	96%
Volta	⬆0.150	97	⬇100	⬇44	1.00	96%
Western	0.040	96	98	⬇43	0.99	89%
Western North	0.040	97	⬆92	47	0.96	97%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2