■ Malaria: Ghana (T4/2019)

Sur la voie
Progrès
Pas sur la voie

N'est pas	applicable
No data	





Indicateurs nationaux

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	100,000 persons	under 5 years old	% of pregnant women who slept under an ITN the previous night		received 3 doses	% of population in target areas protected with IRS in the last 12 months
100%	20.4%	4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Carte de score

Region	In	Impact		Cover	age	
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.110	95	93	↓ 43	0.92	₽ 83%
Ahafo	0.070	98	92	↓ 53	0.97	95%
Ashanti	0.050	92	₽ 90	↓ 41	0.93	₽ 82%
Bono	1 0.050	98	77	65	0.98	91%
Bono East	↓ 0.330	99	1 91	58	1 0.93	1 92%
Central	♦ 0.110	95	91	↓ 45	0.95	88%
Eastern	♦ 0.120	98	↓ 101	45	0.97	90%
Greater Accra	0.320	94	₽ 93	↓ 46	0.89	81%
North East	1 0.000	82	1 49	↓ 17	0.92	1 91%
Northern	↓ 0.190	76	97	↓ 31	0.66	↓ 48%
Oti	0.180	98	102	↓ 35	0.98	99%
Savannah	0.000	80	₽ 94	27	0.94	₽ 82%
Upper East	1 0.000	98	1 00	↓ 48	0.98	1 93%
Upper West	0.120	98	94	↓ 49	0.93	96%
Volta	1 0.150	97	↓ 100	↓ 44	1.00	96%
Western	0.040	96	98	↓ 43	0.99	89%
Western North	0.040	97	1 92	47	0.96	97%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2