Malaria: Ghana (Q1/2020)

On track | Not applicable | ⬆ Improve
Progress | No data | ⬇ Decline
Not on track

Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
33%	20.4%	⬆4		50%	52.6%	59.6%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	⬆0.060	95	96	⬆50	0.90	87%
Ahafo	⬆0.000	98	⬇84	53	0.97	⬆100%
Ashanti	0.040	92	⬆97	42	0.93	84%
Bono	⬇0.110	100	77	⬆106	0.95	92%
Bono East	⬆0.100	99	87	58	0.93	97%
Central	0.150	96	⬆97	⬆51	⬇0.86	⬇81%
Eastern	⬆0.050	99	100	48	0.99	⬆96%
Greater Accra	⬆0.000	94	93	⬆58	0.88	⬆88%
North East	⬇0.110	81	⬆66	18	0.92	91%
Northern	⬆0.070	75	⬆111	29	0.63	⬆63%
Oti	⬆0.000	96	102	36	0.94	⬇94%
Savannah	0.000	78	⬆122	⬇19	0.91	81%
Upper East	0.000	98	100	⬆58	0.97	95%
Upper West	⬆0.000	99	98	⬆56	0.94	93%
Volta	⬆0.000	98	103	⬆52	1.00	93%
Western	⬇0.110	96	98	⬆57	0.96	90%
Western North	0.060	96	⬆97	⬆56	0.93	⬇87%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2