

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	↓0.140	96	96	45	↓0.01	86%
Ahafo	0.000	99	86	53	↓0.01	↓91%
Ashanti	0.070	93	97	42	↓0.01	85%
Bono	↓0.180	99	↑83	↓64	↓0.01	93%
Bono East	↑0.000	97	89	↓47	↓0.01	↓90%
Central	↓0.270	98	↑102	52	↓0.01	↓70%
Eastern	↓0.100	99	102	45	↓0.01	97%
Greater Accra	0.000	96	92	↓50	↓0.01	88%
North East	↓0.200	78	↑75	22	↓0.01	↓85%
Northern	0.100	↑90	107	26	↓0.01	↑73%
Oti	↓0.270	98	106	↑43	↓0.01	92%
Savannah	↓0.770	78	↓98	21	↓0.01	80%
Upper East	0.000	96	98	54	0.95	95%
Upper West	↓0.340	99	↓93	60	↓0.01	89%
Volta	↓0.480	99	104	50	↓0.01	93%
Western	0.070	97	95	54	↓0.01	↓80%
Western North	↓0.160	96	102	↓50	↓0.01	83%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2