



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
↓40%	↑14.0%	↑4		49%	↑67.0%	61.0%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.150	95	97	45	↑0.94	↓36%
Ahafo	↓0.140	100	87	↑60	↑1.00	91%
Ashanti	↓0.160	92	97	↑48	↑0.91	↓79%
Bono	↑0.050	100	82	↑73	↑0.99	95%
Bono East	↓0.450	100	88	50	↑0.96	94%
Central	↑0.170	97	100	49	↑0.93	↑84%
Eastern	0.090	99	101	43	↑0.99	↓7%
Greater Accra	↓0.090	95	↓86	↓43	↑0.94	84%
North East	↓0.280	↓60	↑114	20	↑0.95	86%
Northern	↑0.040	↓82	108	29	↑0.95	↑95%
Oti	↓0.420	97	104	↓36	↑1.00	93%
Savannah	↑0.330	77	↑105	↓16	↑0.96	↑87%
Upper East		99	101	↑67	↑0.96	92%
Upper West	↑0.240	100	90	↑69	↑0.99	↑97%
Volta	↑0.110	99	↓97	↓34	↑1.00	93%
Western	0.070	92	↑106	↓48	↑0.88	79%
Western North	↓0.400	94	↓97	46	↑0.71	↓76%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2