Malaria: Ghana (T3/2020)

Sur la voie
Progrès
Pas sur la voie
N'est pas applicable
No data
Improve
Decline

Indicateurs nationaux

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
⬇40%	⬆14.0%	⬆4		49%	⬆67.0%	61.0%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Carte de score

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.150	95	97	45	⬆0.94	⬇36%
Ahafo	⬇0.140	100	87	⬆60	⬆1.00	91%
Ashanti	⬇0.160	92	97	⬆48	⬆0.91	⬇79%
Bono	⬆0.050	100	82	⬆73	⬆0.99	95%
Bono East	⬇0.450	100	88	50	⬆0.96	94%
Central	⬆0.170	97	100	49	⬆0.93	⬆84%
Eastern	0.090	99	101	43	⬆0.99	⬇7%
Greater Accra	⬇0.090	95	⬇86	⬇43	⬆0.94	84%
North East	⬇0.280	⬇60	⬆114	20	⬆0.95	86%
Northern	⬆0.040	⬇82	108	29	⬆0.95	⬆95%
Oti	⬇0.420	97	104	⬇36	⬆1.00	93%
Savannah	⬆0.330	77	⬆105	⬇16	⬆0.96	⬆87%
Upper East		99	101	⬆67	0.96	92%
Upper West	⬆0.240	100	90	⬆69	⬆0.99	⬆97%
Volta	⬆0.110	99	⬇97	⬇34	⬆1.00	93%
Western	0.070	92	⬆106	⬇48	⬆0.88	79%
Western North	⬇0.400	94	⬇97	46	⬆0.71	⬇76%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2