



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
39%	↑ 14.0%	↑ 4		49%	↑ 67.0%	61.0%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.100	96	94	↓ 40	↑ 1.00	↑ 87%
Ahafo	0.160	98	84	56	↑ 1.10	↑ 98%
Ashanti	↑ 0.020	95	↓ 88	↓ 40	↑ 1.00	84%
Bono	↓ 0.220	99	↓ 73	↓ 62	↑ 1.10	94%
Bono East	↑ 0.210	98	92	↓ 43	↑ 1.10	↑ 99%
Central		97	96	↓ 44	↑ 1.00	87%
Eastern	↓ 0.160	100	99	40	0.96	↑ 92%
Greater Accra	↓ 0.390	95	↑ 93	45	↑ 1.10	82%
North East		62	↑ 121	18	↓ 0.87	↑ 92%
Northern	0.060	↑ 89	↓ 96	25	↑ 1.20	↓ 87%
Oti	↑ 0.170	100	105	↓ 26	↓ 0.95	↑ 99%
Savannah		72	↓ 92	16	0.94	↓ 81%
Upper East		99	97	↓ 51	0.95	92%
Upper West	↑ 0.090	100	92	↓ 45	0.97	93%
Volta	↓ 0.410	99	↑ 103	37	↑ 1.10	94%
Western	0.060	92	102	↓ 39	0.91	77%
Western North	↑ 0.180	97	93	↓ 41	↓ 0.59	72%
Source:	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2