## **■ Malaria: Ghana (T4/2020)**

Sur la voie
Progrès
Pac cur la voio

N'est p	as	appl	icable
No dat	a		





## **Indicateurs nationaux**

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	under 5 years old		% of population using ITN among those with access to an ITN		IRS in the last 12
39%	<b>1</b> 4.0%	<b>1</b> 4		49%	<b>1</b> 67.0%	61.0%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

## Carte de score

Region	In	Impact		Cover	age	
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.100	96	94	<b>↓</b> 40	<b>1</b> .00	<b>↑</b> 87%
Ahafo	0.160	98	84	56	<b>1.10</b>	<b>1</b> 98%
Ashanti	<b>↑</b> 0.020	95	<b>₽</b> 88	<b>↓</b> 40	<b>1</b> .00	84%
Bono	<b>₽</b> 0.220	99	<b>₽</b> 73	<b>↓</b> 62	<b>1.10</b>	94%
Bono East	<b> 0.210</b>	98	92	<b>↓</b> 43	<b>1.10</b>	<b>1</b> 99%
Central		97	96	<b>₽</b> 44	<b>1</b> .00	87%
Eastern	<b>♦</b> 0.160	100	99	40	0.96	<b>1</b> 92%
Greater Accra	<b>♦</b> 0.390	95	<b>1</b> 93	45	<b>1.10</b>	82%
North East		62	<b>1</b> 21	18	<b>↓</b> 0.87	<b>1</b> 92%
Northern	0.060	<b>1</b> 89	<b>₽</b> 96	25	<b>1</b> .20	<b>₽</b> 87%
Oti	<b>↑</b> 0.170	100	105	<b>₽</b> 26	<b>↓</b> 0.95	<b>1</b> 99%
Savannah		72	<b>₽</b> 92	16	0.94	<b>₽</b> 81%
Upper East		99	97	<b>↓</b> 51	0.95	92%
Upper West	<b>1</b> 0.090	100	92	<b>↓</b> 45	0.97	93%
Volta	<b>♦</b> 0.410	99	<b>1</b> 03	37	<b>1.10</b>	94%
Western	0.060	92	102	<b>₽</b> 39	0.91	77%
Western North	<b>1</b> 0.180	97	93	<b>₽</b> 41	<b>₽</b> 0.59	72%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2