



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
34%	↑ 14.0%	4		49%	↑ 67.0%	61.0%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.080	97	97	↑ 51	↓ 0.94	90%
Ahafo	↑ 0.080	98	↑ 92	↑ 67	↓ 0.91	99%
Ashanti	0.030	94	93	↑ 46	0.96	↑ 91%
Bono		99	↑ 79	↑ 67	↓ 1.00	98%
Bono East	↓ 0.550	100	88	44	↓ 0.94	95%
Central	0.110	99	↑ 102	↑ 57	0.99	↑ 96%
Eastern	↑ 0.040	99	101	↑ 54	0.97	92%
Greater Accra		97	90	↑ 61	↓ 0.97	↑ 87%
North East	0.080	↑ 80	122	↓ 12	0.90	↓ 81%
Northern	↓ 0.140	91	92	↑ 38	↓ 0.90	90%
Oti		98	104	↑ 42	0.97	97%
Savannah	0.190	↑ 85	↑ 116	↑ 23	↑ 0.99	↑ 89%
Upper East		99	98	↑ 62	0.99	96%
Upper West	↓ 0.190	100	↑ 101	↑ 62	0.95	↑ 100%
Volta	↑ 0.150	99	↓ 93	↑ 51	↓ 1.00	98%
Western	0.060	92	↑ 109	↑ 52	↓ 0.85	74%
Western North	↑ 0.050	95	98	↑ 57	0.58	↑ 80%
Source:	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2	DHIMS 2