

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.080	97	97	↑51	↓0.94	90%
Ahafo	↑0.080	98	↑92	↑67	↓0.91	99%
Ashanti	0.030	94	93	↑46	0.96	↑91%
Bono		99	↑79	↑67	↓1.00	98%
Bono East	↓0.550	100	88	44	↓0.94	95%
Central	0.110	99	↑102	↑57	0.99	↑96%
Eastern	↑0.040	99	101	↑54	0.97	92%
Greater Accra		97	90	↑61	↓0.97	↑87%
North East	0.080	↑80	122	↓12	0.90	↓81%
Northern	↓0.140	91	92	↑38	↓0.90	90%
Oti		98	104	↑42	0.97	97%
Savannah	0.190	↑85	↑116	↑23	↑0.99	↑89%
Upper East		99	98	↑62	0.99	96%
Upper West	↓0.190	100	↑101	↑62	0.95	↑100%
Volta	↑0.150	99	↓93	↑51	↓1.00	98%
Western	0.060	92	↑109	↑52	↓0.85	74%
Western North	↑0.050	95	98	↑57	0.58	↑80%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2