



Country indicators

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
39%	14.0%	4		49%	67.0%	61.0%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Scorecard

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.050	98	↓5	↑57	0.93	91%
Ahafo		99	↓2	65	↓0.80	↓93%
Ashanti	0.050	96	↓5	↑54	0.94	92%
Bono	0.040	99	↓6	64	1.00	99%
Bono East	↑0.080	99	↓4	↑54	↓0.78	↓84%
Central		100	↓9	↑64	0.99	96%
Eastern	0.070	99	↓6	↑62	0.95	91%
Greater Accra		97	↓6	↑66	↓0.90	91%
North East	↓0.700	↑95	↓10	↑31	↑0.98	↑100%
Northern	↑0.030	95	↓6	↑54	0.86	↓82%
Oti	0.230	97	↓6	↑53	1.00	93%
Savannah		↑91	↓5	27	↓0.92	86%
Upper East		99	↓5	↓44	1.00	↓89%
Upper West	↑0.090	100	↓2	65	0.98	98%
Volta		98	↓10	↑59	0.99	96%
Western	↓0.110	96	↓3	↑59	↑0.90	↑87%
Western North	↓0.110	97	↓1	↑66	↑0.78	↑86%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2