Malaria: Ghana (T2/2021)

Sur la voie
Progrès
Pas sur la voie
N'est pas applicable
No data
Improve
Decline

Indicateurs nationaux

Malaria test positivity ratio (sentinel sites)	% Parasitemia prevalence: children aged 6–59 months with malaria infection (by microscopy)	Inpatient malaria deaths per 100,000 persons per year	% of children under 5 years old who slept under an ITN the previous night	% of pregnant women who slept under an ITN the previous night	% of population using ITN among those with access to an ITN	% of pregnant women who received 3 doses of IPT for malaria during ANC visits during their last pregnancy	% of population in target areas protected with IRS in the last 12 months
39%	14.0%	4		49%	67.0%	61.0%	84.9%
DHIMS 2	Survey	DHIMS 2	Survey	Survey	Surveys	Surveys	IRS report

Carte de score

Region	Impact		Coverage			
	Under five case fatality rate	% of suspected malaria cases that received a parasitological test (RDTs or microscopy)	% of confirmed malaria cases treated with ACT at health facilities	% of pregnant women on Intermittent preventive treatment at least three doses at ANC according to national policy	Proportion of ANC registrants given ITN	Proportion of children due for measles 2 given ITN
Ghana	0.050	98	⬇5	⬆57	0.93	91%
Ahafo		99	⬇2	65	⬇0.80	⬇93%
Ashanti	0.050	96	⬇5	⬆54	0.94	92%
Bono	0.040	99	⬇6	64	1.00	99%
Bono East	⬆0.080	99	⬇4	⬆54	⬇0.78	⬇84%
Central		100	⬇9	⬆64	0.99	96%
Eastern	0.070	99	⬇6	⬆62	0.95	91%
Greater Accra		97	⬇6	⬆66	⬇0.90	91%
North East	⬇0.700	⬆95	⬇10	⬆31	⬆0.98	⬆100%
Northern	⬆0.030	95	⬇6	⬆54	0.86	⬇82%
Oti	0.230	97	⬇6	⬆53	1.00	93%
Savannah		⬆91	⬇5	27	⬇0.92	86%
Upper East		99	⬇5	⬇44	1.00	⬇89%
Upper West	⬆0.090	100	⬇2	65	0.98	98%
Volta		98	⬇10	⬆59	0.99	96%
Western	⬇0.110	96	⬇3	⬆59	⬆0.90	⬆87%
Western North	⬇0.110	97	⬇1	⬆66	⬆0.78	⬆86%
Source:	DHIMS 2	DHIMS 2	DHMIS 2	DHIMS 2	DHIMS 2	DHIMS 2