

Indicateurs nationaux

% HfTs that attained UC of LLINs in targeted areas	% PW who slept under an LLIN among HfTs with at least 1 net	% of pregnant women who received 3+ doses of IPTp for malaria during their last pregnancy in the last 2 yrs	% HfTs having no stock-out of ACTs	ACTs distributed against projected Forecasting & Quantification targets	% patients treated according to national guidelines	Malaria prevalence in children <5 yrs		
51%	85%	83%	38%	62%	84%	100%	94%	5%
DHS/KMIS	DHS/KMIS / DHS/KMIS	KMIS	QOC survey / QOC Survey	KEMSA	QOC Survey	KMIS		

Carte de score

Region	Malaria Prevention			Malaria in Pregnancy			Case Management			EPR	Surveillance Monitoring & Evaluation and Operational Research			
	Transmission Zone	% PW receiving an LLIN at ANC clinic in targeted areas	% children <1 yr receiving an LLIN at immunization clinic in targeted areas	% PW who received IPT 2 in ANC malaria endemic areas	% PW who had 4 ANC visits	% suspected malaria cases receiving a parasitological test	Annual Blood Examination Rate	Malaria cases per 1000 population	Outpatient cases treated w/ AL as a proportion of confirmed malaria cases		% epidemic prone and seasonal transmission sub counties with updated annual EPR plans	Reporting rates of MOH S05 (weekly surveillance)	Reporting rate for malaria commodity form	Annual entomological inoculation rate / Confirmed Malaria Cases per 1,000 Population per Year
Kenya		55	61	53	49	69%	2%	16	103	100%	76%	93%	4.29	
Baringo	Seasonal, arid, & semi-arid	22	30	0	40	57%	1%	15	132	100%	49%	95%	0.83	
Bomet	Highland epidemic prone	18	26	0	39	79%	0%	0	178	100%	74%	100%	0	
Bungoma	Lake endemic	86	85	64	53	82%	9%	35	86	100%	82%	100%	0	
Busia	Lake endemic	76	82	73	49	80%	14%	80	86	0%	96%	94%	33.97	
Elgeyo-Marakwet	Seasonal, arid, & semi-arid	14	11	0	30	100%	1%	9	172	100%	77%	90%	0	
Embu	Seasonal, arid, & semi-arid	46	67	0	45	100%	0%	0	57	100%	78%	100%	0	
Garissa	Seasonal, arid, & semi-arid	0	0	0	61	100%	0%	1	28	100%	95%	92%	0	
Homa Bay	Lake endemic	81	81	37	43	64%	3%	22	107	0%	86%	97%	7.32	
Isiolo	Seasonal, arid, & semi-arid	28	37	0	42	100%	1%	3	35	100%	61%	65%	0	
Kajiado	Seasonal, arid, & semi-arid	73	80	0	53	52%	0%	1	27	100%	92%	96%	0	
Kakamega	Lake endemic	67	64	49	55	64%	8%	57	99	100%	100%	100%	6.33	
Kericho	Highland epidemic prone	9	14	0	37	100%	1%	3	89	100%	93%	82%	0.99	
Kiambu	Low risk	0	0	0	52	100%	0%	0	25	0%	56%	85%	0	
Kilifi	Coast endemic	121	105	62	51	38%	1%	31	85	0%	68%	95%	1.81	
Kirinyaga	Low risk	8	17	0	56	44%	0%	0	25	0%	77%	100%	3.01	
Kisii	Highland epidemic prone	9	12	0	44	62%	2%	6	94	100%	93%	90%	9.94	
Kisumu	Lake endemic	77	85	64	58	78%	12%	49	143	0%	78%	100%	13.86	
Kitui	Seasonal, arid, & semi-arid	0	0	0	31	39%	0%	0	68	100%	93%	88%	0	
Kwale	Coast endemic	101	96	55	46	12%	1%	25	210	0%	35%	75%	0.8	
Lalikipia	Low risk	0	0	0	70	49%	0%	0	16	0%	59%	94%	0	
Lamu	Coast endemic	129	123	66	57	62%	0%	1	162	0%	75%	94%	0	
Machakos County	Low risk	16	25	0	54	94%	0%	0	9	0%	78%	90%	0	
Makueni	Low risk	48	81	0	60	42%	0%	0	50	0%	100%	94%	0	
Mandera	Seasonal, arid, & semi-arid	0	0	0	40	19%	0%	0	64	100%	39%	56%	0	
Marsabit	Seasonal, arid, & semi-arid	0	0	0	49	49%	0%	1	54	100%	55%	94%	0	
Meru	Seasonal, arid, & semi-arid	47	74	0	37	28%	0%	1	51	100%	72%	86%	0	
Migori	Lake endemic	78	84	52	51	77%	8%	40	102	0%	96%	100%	14.25	
Mombasa	Coast endemic	101	82	49	36	62%	1%	5	16	0%	57%	95%	0	
Muranga	Low risk	42	63	0	47	69%	0%	0	30	0%	61%	87%	0	
Nairobi	Low risk	0	0	0	58	91%	0%	2	16	0%	50%	91%	0	
Nakuru	Low risk	0	0	0	53	45%	0%	1	36	0%	77%	97%	0	
Nandi	Highland epidemic prone	42	42	0	35	67%	1%	6	156	100%	71%	78%	3.35	
Narok	Highland epidemic prone	37	45	0	38	35%	0%	2	236	100%	69%	93%	0	
Nyamira	Highland epidemic prone	42	47	0	72	47%	0%	4	47	100%	71%	99%	1.04	
Nyandarua	Low risk	0	0	0	42	100%	0%	0	10	0%	61%	98%	0	
Nyeri	Low risk	0	0	0	65	59%	0%	0	8	0%	78%	98%	0	
Samburu	Seasonal, arid, & semi-arid	0	0	0	37	53%	0%	6	92	100%	46%	92%	0	
Siaya	Lake endemic	90	98	58	57	70%	13%	104	104	0%	74%	96%	46.66	
Taita Taveta	Coast endemic	119	98	70	54	100%	1%	1	55	0%	66%	74%	0.88	
Tana River	Seasonal, arid, & semi-arid	77	54	40	26	26%	0%	3	118	100%	46%	92%	0	
Tharaka Nithi	Seasonal, arid, & semi-arid	59	67	0	63	79%	1%	2	44	100%	100%	100%	0	
Trans-Nzoia	Highland epidemic prone	47	65	0	44	74%	1%	16	69	100%	74%	91%	1.76	
Turkana	Seasonal, arid, & semi-arid	29	0	0	53	100%	5%	115	94	100%	88%	93%	2.59	
Uasin Gishu	Highland epidemic prone	57	76	1	48	100%	1%	3	51	100%	87%	97%	0	
Vihiga	Lake endemic	68	74	57	49	81%	8%	35	88	0%	100%	100%	0.86	
Wajir	Seasonal, arid, & semi-arid	0	0	0	33	36%	0%	0	165	100%	84%	89%	0	
West Pokot	Seasonal, arid, & semi-arid	5	6	0	27	72%	3%	39	157	100%	90%	96%	0	
Source:	Policy	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	NMCP	DHIS	DHIS	Activity Report / DHIS	