

Country indicators

% HHS that have attained UC of LLINs in targeted areas	% PW who slept under an LLIN among HHS with at least 1 net	% of pregnant women who received 3+ doses of IPTp for malaria during their last pregnancy in the last 2 yrs	% HHS having no stock-out of ACTs	ACTs distributed against projected Forecasting & Quantification targets	% patients treated according to national guidelines	Malaria prevalence in children <5 yrs
51%	85%	83%	38%	71%	81%	
DHS/KMIS	DHS/KMIS / DHS/KMIS	KMIS	QOC survey / QOC Survey	KEMSA	QOC Survey	KMIS

Scorecard

Region	Malaria Prevention			Malaria in Pregnancy			Case Management			EPR	Surveillance Monitoring & Evaluation and Operational Research			
	Transmission Zone	% PW receiving an LLIN at ANC clinic in targeted areas	% children <1 yr receiving an LLIN at immunization clinic in targeted areas	% PW who receiving IPT 2 in ANC malaria endemic areas	% PW who had 4 ANC visits	% suspected malaria cases receiving a parasitological test	Annual Blood Examination Rate	Malaria cases per 1000 population	Outpatient cases treated w/AL as a proportion of confirmed malaria cases		% epidemic prone and seasonal transmission sub counties with updated annual EPR plans	Reporting rates of MOH S05 (weekly surveillance)	Reporting rate for malaria commodity form	Annual entomological inoculation rate / Confirmed Malaria Cases per 1,000 Population per Year
Kenya		86	78	57	44	74%	8%	17	89	100%	69%	85%	4.29	
Baringo	Seasonal, arid, & semi-arid	69	89	0	31	46%	2%	12	106	100%	39%	70%	0.83	
Bomet	Highland epidemic prone	81	106	0	42	50%	1%	1	157	100%	82%	87%	0	
Bungoma	Lake endemic	96	83	79	45	86%	31%	50	75	100%	77%	100%	0	
Busia	Lake endemic	108	83	41	33	91%	49%	75	71	0%	89%	100%	33.97	
Elgeyo-Marakwet	Seasonal, arid, & semi-arid	86	98	0	24	100%	0%	6	48	100%	76%	92%	0	
Embu	Seasonal, arid, & semi-arid	83	93	0	43	100%	0%	0	52	100%	63%	94%	0	
Garissa	Seasonal, arid, & semi-arid	0	0	0	45	74%	1%	1	27	100%	76%	74%	0	
Homa Bay	Lake endemic	84	63	52	39	43%	9%	28	101	0%	78%	91%	7.32	
Isiolo	Seasonal, arid, & semi-arid	143	167	0	54	76%	1%	2	91	100%	63%	84%	0	
Kajiado	Seasonal, arid, & semi-arid	57	50	0	48	33%	1%	1	15	100%	83%	91%	0	
Kakamega	Lake endemic	97	87	60	44	78%	45%	68	101	100%	92%	100%	6.33	
Kericho	Highland epidemic prone	47	55	0	31	92%	5%	5	74	100%	84%	65%	0.99	
Kiambu	Low risk	0	0	0	48	100%	0%	1	17	0%	62%	80%	0	
Kilifi	Coast endemic	107	61	60	44	46%	5%	16	191	0%	51%	92%	1.81	
Kirinyaga	Low risk	0	0	0	53	100%	1%	0	72	0%	65%	93%	3.01	
Kisii	Highland epidemic prone	78	76	0	28	66%	5%	7	64	100%	88%	90%	9.94	
Kisumu	Lake endemic	68	61	49	49	82%	19%	71	71	0%	54%	100%	13.86	
Kitui	Seasonal, arid, & semi-arid	82	59	0	36	38%	1%	1	39	100%	87%	91%	0	
Kwale	Coast endemic	84	65	63	33	40%	0%	34	129	0%	12%	80%	0.8	
Laikipia	Low risk	0	0	0	57	52%	0%	0	34	0%	60%	63%	0	
Lamu	Coast endemic	97	70	46	57	45%	0%	1	42	0%	74%	77%	0	
Machakos County	Low risk	104	118	0	52	87%	1%	1	14	0%	80%	86%	0	
Makueni	Low risk	100	125	0	63	41%	0%	1	12	0%	92%	97%	0	
Mandera	Seasonal, arid, & semi-arid	0	0	0	38	16%	0%	0	260	100%	18%	39%	0	
Marsabit	Seasonal, arid, & semi-arid	0	0	0	48	34%	1%	1	22	100%	34%	88%	0	
Meru	Seasonal, arid, & semi-arid	0	0	0	37	27%	1%	1	56	100%	61%	60%	0	
Migori	Lake endemic	67	54	25	35	80%	37%	63	79	0%	88%	100%	14.25	
Mombasa	Coast endemic	88	46	71	46	61%	10%	6	7	0%	65%	72%	0	
Muranga	Low risk	71	61	0	49	41%	0%	0	13	0%	66%	66%	0	
Nairobi	Low risk	0	0	0	58	80%	1%	2	14	0%	39%	66%	0	
Nakuru	Low risk	0	0	0	55	57%	1%	2	22	0%	81%	77%	0	
Nandi	Highland epidemic prone	22	30	0	29	69%	4%	9	104	100%	64%	85%	3.35	
Narok	Highland epidemic prone	66	80	0	34	43%	6%	4	135	100%	68%	88%	0	
Nyamira	Highland epidemic prone	77	68	0	56	52%	0%	5	54	100%	65%	75%	1.04	
Nyandarua	Low risk	0	0	0	38	46%	0%	0	7	0%	55%	91%	0	
Nyeri	Low risk	0	0	0	57	31%	0%	0	12	0%	75%	76%	0	
Samburu	Seasonal, arid, & semi-arid	0	0	0	42	35%	0%	5	75	100%	54%	95%	0	
Siaya	Lake endemic	93	81	60	51	70%	33%	106	105	0%	63%	98%	46.66	
Taita Taveta	Coast endemic	49	26	35	24	100%	0%	1	55	0%	59%	52%	0.88	
Tana River	Seasonal, arid, & semi-arid	115	84	61	52	38%	5%	6	87	100%	41%	79%	0	
Tharaka Nithi	Seasonal, arid, & semi-arid	103	88	0	62	54%	1%	4	1	100%	89%	87%	0	
Trans-Nzoia	Highland epidemic prone	57	52	0	25	43%	2%	16	45	100%	66%	81%	1.76	
Turkana	Seasonal, arid, & semi-arid	0	0	0	54	65%	9%	45	99	100%	83%	89%	2.59	
Uasin Gishu	Highland epidemic prone	98	88	0	42	82%	2%	4	81	100%	84%	93%	0	
Vihiga	Lake endemic	174	126	89	39	81%	23%	44	67	0%	85%	98%	0.86	
Wajir	Seasonal, arid, & semi-arid	0	0	0	36	53%	0%	0	171	100%	75%	73%	0	
West Pokot	Seasonal, arid, & semi-arid	0	0	0	26	46%	4%	17	109	100%	74%	94%	0	
Source:	Policy	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	NMCP	DHIS	DHIS	Activity Report / DHIS	