■ Malaria Scorecard: Kenya (T1/2021)

Indicateurs nationaux

% HHs that have attained UC of LLINs in targeted areas	% PW who slept under an LLIN among HHs with at least 1 net	% of pregnant women who received 3+ doses of IPTp for malaria during their last pregnancy in the last 2 yrs		ACTs distributed against projected Forecasting & Quantification targets	% patients treated according to national guidelines	Malaria prevalence in children <5 yrs	
51%	85% 83%	38%	•71% 81%		92%	5%	
DHS/KMIS	DHS/KMIS / DHS/KMIS	KMIS	QOC survey / QOC Survey	KEMSA	QOC Survey	KMIS	

Carte de score

Region	Malaria Prevention			Malaria in Pregnancy		Case Management				EPR	Surveillance Monitoring & Evaluation and Operational Research		
	Transmission Zone	an LLIN at ANC	% children <1 yr receiving an LLIN at immunization clinic in targeted areas	in ANC malaria		% suspected malaria cases receiving a parasitological test	Annual Blood Examination Rate	Malaria cases per 1000 population	Outpatient cases treated w/ AL as a proportion of confirmed malaria cases	% epidemic prone and seasonal transmission sub counties with updated annual EPR plans	Reporting rates of	Reporting rate	Annual entomologica
Kenya		* 86	1 78	57	44	74%	18%	17	♦ 89	100%	 69%	+ 85%	4.29
Baringo County	Seasonal, arid, & semi-arid	1 69	1 89	0	+31	4 46%	◆ 2%	12	♦ 106	100%	♦ 39%	₹ 70%	0.83
Bomet County	Highland epidemic prone	± 81	1 06	0	42	₹ 50%	± 196	1	♦ 157	100%	± 82%	♦ 87%	0
Bungoma County	Lake endemic	1 96	83	1 79	₽ 45	86%	± 31%	♦ 50	♦75	100%	₹77%	100%	0 16
Busia County	Lake endemic	± 108	83	♦ 41	+33	± 91%	± 49%	75	◆71	0%	♦89%	± 100%	33.97
Elgeyo Marakwet County	Seasonal, arid, & semi-arid	± 86	1 98	0	₽ 24	100%	₽ 0%	6	♦ 48	100%	76%	92%	0
Embu County	Seasonal, arid, & semi-arid	± 83	1 93	0	43	100%	0%	0	♦ 52	100%	4 63%	494%	0
Garissa County	Seasonal, arid, & semi-arid	0	0	0	₽ 45	74%	196	1		100%	₹ 76%	₹ 74%	0
Homa Bay County	Lake endemic	84	4 63	± 52	39	4 43%	* 9%	4 28	♦ 101	0%	₹ 78%	 91%	7.32
Isiolo County	Seasonal, arid, & semi-arid	1 143	1 67	0	± 54	76%	196	2	± 91	100%	63%	* 84%	0
Kajiado County	Seasonal, arid, & semi-arid	\$ 57	\$ 50	0	 48	\$ 33%	196	1	 15	100%	₽ 83%	91%	0
Kakamega County	Lake endemic	•97	1 87	1 60	₽ 44	↑ 78%	1 45%	4 68		100%	♦ 92%	100%	6.33
Kericho County	Highland epidemic prone	1 47	1 55	0	\$ 31	92%	+ 5%	5	♦ 74	100%	♦ 84%	♦ 65%	0.99
Kiambu County	Low risk	0	0	0	48	100%	0%	1	♦ 17	0%	1 62%	80%	0
Kilifi County	Coast endemic	♦107	♦ 61	60	♣ 44	* 46%	1 5%	1 16	1 91	0%	♦ 51%	92%	1.81
Kirinyaga County	Low risk	0	0	0	53	100%	1 96	0	+ 72	0%	◆ 65%	♦ 93%	3.01
Kisii County	Highland epidemic prone	★ 78	+ 76	0	♦ 28	66%	★ 596	7	♣ 64	100%	88%	90%	9.94
Kisumu County	Lake endemic	4 68	♦ 61	44 9	♦ 49	82%	1 9%	₽ 71	₹ 71	0%	♦ 54%	100%	13.86
Kitui County	Seasonal, arid, & semi-arid	82	59	0	36	38%	1 96	1	♦ 39	100%	₽ 87%	91%	0
Kwale County	Coast endemic	₽ 84	4 65	 6 3	4 33	± 40%	. 0%	≢34	1 129	0%	1 2%	* 80%	0.8
Laikipia County	Low risk	0	0	0	◆ 57	52%	0%	0	+ 34	0%	60%	•63% •63%	0.8
		4 97	\$ 70	♦ 46	57	\$45%		•	+ 42	0%	74%	₹77%	0
Lamu County Machakos County	Coast endemic Low risk	◆97 ◆104	◆ 70 • 118	0	52	+ 45% + 87%	1%		+ 42 1 14	0%	74%	86%	0
Makueni County	Low risk	1 04 1 00	1 110 125	0	63	41%	0%		* 14 * 12	0%	•92%	97%	0
· · · · · · · · · · · · · · · · · · ·		0	0	0	38	16%	0%	,	◆12 ◆260		◆92% ◆18%		0
Mandera County	Seasonal, arid, & semi-arid			-	48	± 34%	196			100%	◆18% ◆34%	♦ 39%	0
Marsabit County	Seasonal, arid, & semi-arid	. 0	0	0	37				◆22 56			+ 88% + 60%	
Meru County	Seasonal, arid, & semi-arid	0	0	0		27%	1 96			100%	ቆ 61%	♦ 60%	14.25
Migori County	Lake endemic	♦ 67	♦ 54	♦ 25	♦ 35	80%	* 37%	♦ 63	♦ 79	0%	♦88%	100%	0 +20
Mombasa County	Coast endemic	♦ 88	. 46	◆71	± 46	61%	1 0%	6	. 7	0%	* 65%	₹ 72%	0 2
Muranga County	Low risk	± 71	61	0	49	. 41%	0%		 13	0%	66%	4 66%	0
Nairobi County	Low risk	0	0	0	58	\$ 80%	1 96	2	14	0%	4 39%	. 66%	0
Nakuru County	Low risk	0	0	0	55	1 57%	1 96	2	\$ 22	0%	81%	\$ 77%	3.35
Nandi County	Highland epidemic prone	1 22	1 30	. 0	\$ 29	. 69%	1 4%	9	+ 104	100%	₽ 64%	1 85%	0
Narok County	Highland epidemic prone	• 66	1 80	0	34	1 43%	1 6%	4	1 35	100%	68%	\$ 88%	
Nyamira County	Highland epidemic prone	1 77	◆ 68	0	₹ 56	52%	0%	5	1 54	100%	₹ 65%	₹ 75%	1.04
Nyandarua County	Low risk	. 0	0	0	38	46%	0%			0%	₹55%	≠ 91%	0
Nyeri County	Low risk	. 0	0	0	♦ 57	♦31%	096	0		0%	75%	₹ 76%	0
Samburu County	Seasonal, arid, & semi-arid	0	0	0	42	♦35 %	0%	5	₹ 75	100%	1 54%	95%	0 •2
Siaya County	Lake endemic	93	₽ 81	60	₽ 51	70%	± 33%	106	105	0%	♦ 63%	98%	46.66
Taita Taveta County	Coast endemic	4 49	♦ 26	♦ 35	₽ 24	100%	◆ 0%	1		0%	◆ 59%	♦52%	0.88
Tana River County	Seasonal, arid, & semi-arid	+ 115	± 84	± 61	+ 52	± 38%	* 5%	6	♦ 87	100%	4 4196	₹ 79%	01
Tharaka Nithi County	Seasonal, arid, & semi-arid	± 103	1 88	0	62	\$ 54%	1 96	4	4 1	100%	₽ 89%	\$ 87%	0
Trans Nzoia County	Highland epidemic prone	• 57	\$ 52	0	\$ 25	4 43%	* 2%	16	4 45	100%	 466%	\$ 81%	1.76
Turkana County	Seasonal, arid, & semi-arid	0	0	0	54	65%	1 9%	1 45	99	100%	83%	89%	2.59
Uasin Gishu County	Highland epidemic prone	● 98	1 88	0	+ 42	82%	1 2%	4	1 81	100%	84%	93%	0 1
Vihiga County	Lake endemic	174	1 26	1 89	♦ 39	81%	1 23%	4 44	4 67	0%	♦85%	98%	0.86
Wajir County	Seasonal, arid, & semi-arid	0	0	0	36	* 53%	0%	0	171	100%	₹ 75%	₹73%	0
West Pokot County	Seasonal, arid, & semi-arid	0	0	0	26	♦ 46%	◆ 4%	1 7	♦ 109	100%	₹ 74%	94%	0 13
Source:	Policy	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	NMCP	DHIS	DHIS	Activity Report / DHIS