

## Country indicators

| % HHs<br>that have<br>attained<br>UC of<br>LLINs in<br>targeted<br>areas | % PW who slept under<br>an LLIN among HHs<br>with at least 1 net | % of pregnant<br>women who<br>received 3+<br>doses of IPTp for<br>malaria during<br>their last<br>pregnancy in the<br>last 2 yrs |                                  | ACTs distributed<br>against projected<br>Forecasting &<br>Quantification<br>targets | % patients<br>treated according<br>to national<br>guidelines | Malaria<br>prevalence in<br>children <5 yrs |  |
|--|--|--|----------------------------------|---|--|---|--|
| 51%  | 85% 83%  | 38%  | •71%<br>81%                      | 100%  | 93%  | 5%  |  |
| DHS/KMIS   | DHS/KMIS / DHS/KMIS  | KMIS   | QOC<br>survey /<br>QOC<br>Survey | KEMSA   | QOC Survey   | KMIS  |  |
| Score  | oord   |  |                                  |   |  |   |  |

| Region                 | Malaria Prevention          |   |   | Malaria in Pregnancy |                              | Case Management  |                                     |                                      |   | EPR   |  | Surveillance Monitoring & Evaluation and<br>Operational Research |  |  |
|------------------------|-----------------------------|---|---|----------------------|------------------------------|--|-------------------------------------|--------------------------------------|---|---|--|--|--|--|
|                        | Transmission Zone           | % PW receiving<br>an LLIN at ANC<br>clinic in targeted<br>areas | % children <1 yr<br>receiving an LLIN<br>at immunization<br>clinic in targeted<br>areas | in ANC malaria       | % PW who had 4<br>ANC visits | % suspected<br>malaria cases<br>receiving a<br>parasitological<br>test | Annual Blood<br>Examination<br>Rate | Malaria cases per<br>1000 population | Outpatient cases<br>treated w/ AL as<br>a proportion of<br>confirmed<br>malaria cases | % epidemic prone and<br>seasonal transmission<br>sub counties with<br>updated annual EPR<br>plans | Reporting rates of<br>MOH 505 (weekly<br>surveillance) | for malaria  | Annual<br>entomological<br>innoculation rat<br>/ Confirmed<br>Malaria Cases<br>per 1,000<br>Population per<br>Year |  |
| Kenya                  |                             | <b>■</b> 62   | <b>₽</b> 63   | <b>*</b> 68          | <b>\$</b> 37                 | 77%  | 1196                                | <b>\$</b> 26                         | 87  | 100%  | <b>*</b> 76%   | <b>*</b> 93%   | 4.29   |  |
| Baringo County         | Seasonal, arid, & semi-arid | 0   | 0   | 0                    |                              | 46%  | <b>*</b> 3%                         | 11                                   | 123   | 100%  | €52%   | <b>●</b> 89%   | 0.83   |  |
| Bomet County           | Highland epidemic prone     | ♦69   | <b>₽</b> 84   | 0                    | <b>#</b> 29                  | <b>\$</b> 38%  | 1%                                  | 1                                    | 159   | 100%  | 79%  | 85%  | 0  |  |
| Bungoma County         | Lake endemic                | ₹74   | €72   | 78                   | 43                           | 83%  | <b>\$</b> 41%                       | ♦85                                  | €85   | 100%  | <b>●</b> 91%   | 100%   | 0 16   |  |
| Busia County           | Lake endemic                | ♦81   | 81  | <b>*</b> 104         | <b>*</b> 39                  | 91%  | <b>*</b> 86%                        | <b>₽</b> 152                         | €84   | 0%  | <b>*</b> 95%   | 100%   | 33.97  |  |
| Elgeyo Marakwet County | Seasonal, arid, & semi-arid | 0   | 0   | 0                    |                              | 88%  | 0%                                  | 6                                    | <b>*</b> 57   | 100%  | €88%   | <b>\$</b> 86%  | 0 +3   |  |
| Embu County            | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | <b>\$</b> 37                 | 98%  | 0%                                  | 0                                    | <b>€</b> 62   | 100%  | ↑75%   | 99%  | 0  |  |
| Garissa County         | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | <b>€</b> 67                  | <b>€</b> 63%   | 1%                                  | 2                                    | <b>●</b> 37   | 100%  | 77%  | <b>●</b> 93%   | 0  |  |
| Homa Bay County        | Lake endemic                | ♦62   | 59  | <b>*</b> 64          | 36                           | <b>*</b> 57%   | <b>*</b> 10%                        | 32                                   | <b>₽</b> 23   | 0%  | 81%  | <b>1</b> 00%   | 7.32 9   |  |
| Isiolo County          | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | <b>\$</b> 35                 | <b>\$</b> 57%  | 1%                                  | 4                                    | <b>₽</b> 84   | 100%  | 62%  | ¥76%   | 0 7 +1   |  |
| Kajiado County         | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | <b>\$</b> 37                 | 37%  | <b>+</b> 0%                         | 1                                    | <b>\$</b> 55  | 100%  | 85%  | 93%  | 0 7.   |  |
| Kakamega County        | Lake endemic                | <b>1</b> 06   | <b>1</b> 02   | <b>€</b> 69          | 41                           | <b>1</b> 83%   | <b>↑</b> 77%                        | <b>₽</b> 136                         | 99  | 100%  | <b>1</b> 99%   | 100%   | 6.33   |  |
| Kericho County         | Highland epidemic prone     | €64   | 57  | 0                    | 27                           | ♦77%   | 5%                                  | 8                                    | €79   | 100%  | €96%   | <b>*</b> 96%   | 0.99   |  |
| Kiambu County          | Low risk                    | 0   | 0   | 0                    | <b></b> 41                   | 63%  | 0%                                  | 0                                    | <b>\$</b> 27  | 0%  | <b>\$</b> 80%  | <b>*</b> 88%   | 0  |  |
| Kilifi County          | Coast endemic               | ₹76   | <b>*</b> 68   | ♦52                  | <b>\$</b> 35                 | <b>\$</b> 38%  | <b>*</b> 7%                         | 13                                   | ♦81   | 0%  | 51%  | 96%  | 1.81   |  |
| Kirinyaga County       | Low risk                    | 0   | 0   | 0                    | <b>\$</b> 42                 | 73%  | 1%                                  | 0                                    | <b>\$</b> 59  | 0%  | <b>1</b> 72%   | 98%  | 3.01   |  |
| Kisii County           | Highland epidemic prone     | <b>€</b> 63   | ♦68   | 0                    | 30                           | <b>↑</b> 74%   | <b>↑</b> 7%                         |                                      | €76   | 100%  | <b>1</b> 99%   | <ul> <li>● 96%</li> </ul>  | 9.94 2   |  |
| Kisumu County          | Lake endemic                | <b>*</b> 95   | <b>1</b> 94   | <b>€</b> 63          | 45                           | 83%  | <b>€</b> 31%                        | ♦87                                  | <b>1</b> 91   | 0%  | 52%  | ♦92%   | 13.86  |  |
| Kitui County           | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | 33                           | <b>♦</b> 30%   | 1%                                  | 0                                    | <b>*</b> 50   | 100%  | <b>*</b> 92%   | 91%  | 0  |  |
| Kwale County           | Coast endemic               | ₹78   | <b>*</b> 79   | <b>*</b> 70          |                              | <b>\$</b> 27%  | <b>*</b> 4%                         | 37                                   | <b>+</b> 120  | 0%  | <b>*</b> 31%   | <b>*</b> 96%   | 0.8  |  |
| Laikipia County        | Low risk                    | 0   | 0   | 0                    | <b>\$</b> 38                 | <b>*</b> 58%   | 0%                                  | 1                                    | <b>+</b> 17   | 0%  | <b>1</b> 71%   | <b>*</b> 99%   | 0  |  |
| Lamu County            | Coast endemic               | 96  | <b>*</b> 82   | <b>●</b> 90          | ♦46                          | <b>♦</b> 37%   | 0%                                  | 0                                    | <b>*</b> 60   | 0%  | <b></b> €15%   | <b></b> €97%   | 0  |  |
| Machakos County        | Low risk                    | 0   | 0   | 0                    | €45                          | <b>1</b> 00%   | <b>♦</b> 0%                         | 0                                    | <b>*</b> 100  | 0%  | <b>1</b> 00%   | <b>*</b> 98%   | 0  |  |
| Makueni County         | Low risk                    | 0   | 0   | 0                    | ₹51                          | <b>₽</b> 26%   | <b>*</b> 4%                         | 0                                    | <b>1</b> 7  | 0%  | <b>*</b> 100%  | 100%   | 0 7  |  |
| Mandera County         | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | <b>\$</b> 30                 | <b>♦</b> 0%  | 0%                                  | 0                                    | <b>#</b> 76   | 100%  | <b>4</b> 0%  | <b>*</b> 67%   | 0  |  |
| Marsabit County        | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | +33                          | <b>↑</b> 76%   | +0%                                 | 2                                    | <b>*</b> 34   | 100%  | <b>↓</b> 27%   | 87%  | 0  |  |
| Meru County            | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | <b>₽</b> 22                  | <b>♦</b> 0%  | <b>₽</b> 0%                         | 0                                    | <b>€</b> 65   | 100%  | <b>♦</b> 37%   | <b>♦</b> 31%   | 0  |  |
| Migori County          | Lake endemic                | <b>*</b> 98   | <b>*</b> 92   | <b>€</b> 63          |                              | 79%  | <b>1</b> 38%                        | ♦80                                  | <b>*</b> 86   | 0%  | 92%  | 100%   | 14.25  |  |
| Mombasa County         | Coast endemic               | • 59  | <b>*</b> 61   | 76                   | <b>\$</b> 37                 | \$54%  | <b>4</b> 7%                         | 4                                    | <b>*</b> 23   | 0%  | <b>*</b> 77%   | <b>*</b> 94%   | 0  |  |
| Muranga County         | Low risk                    | 0   | 0   | 0                    | <b>\$</b> 39                 | <b>*</b> 56%   | 0%                                  | 0                                    |   | 0%  | 62%  | <b>*</b> 96%   | 0  |  |
| Nairobi County         | Low risk                    | 0   | 0   | 0                    | ♦45                          | ♦66%   | 196                                 | 2                                    | <b>€</b> 28   | 0%  | <b>1</b> 51%   | <b>€</b> 87%   | 0 7  |  |
| Nakuru County          | Low risk                    | 0   | 0   | 0                    | <b>₽</b> 36                  | 61%  | <b>₽</b> 0%                         | 2                                    | <b>*</b> 34   | 0%  | <b>₽</b> 75%   | <b>€</b> 85%   | • 7.   |  |
| Nandi County           | Highland epidemic prone     | <b>*</b> 113  | <b>*</b> 150  | 0                    | 29                           | <b>1</b> 79%   | <b>1</b> 6%                         | ♦16                                  | <b>♦</b> 93   | 100%  | <b>*</b> 70%   | <b>1</b> 91%   | 3.35   |  |
| Narok County           | Highland epidemic prone     | <b>*</b> 85   | <b>*</b> 98   | 0                    | <b>₽</b> 27                  | <b>■</b> 31%   | <b>↓</b> 5%                         | 4                                    | 132   | 100%  | <b>*</b> 77%   | <b>•</b> 78%   | 0  |  |
| Nyamira County         | Highland epidemic prone     | 76  | 71  | 0                    | 51                           | <b>\$</b> 43%  | <b>€</b> 2%                         | 10                                   | <b>*</b> 60   | 100%  | <b>1</b> 74%   | <b>*</b> 97%   | 1.04   |  |
| Nyandarua County       | Low risk                    | 0   | 0   | 0                    | 35                           | 100%   | 0%                                  | 0                                    | •19   | 0%  | <b>1</b> 69%   | <b>1</b> 00%   | 0  |  |
| Nyeri County           | Low risk                    | 0   | 0   | 0                    | <b>€</b> 42                  | 27%  | 0%                                  | 0                                    | <b>•</b> 52   | 0%  | 77%  | <b>1</b> 00%   | ° –  |  |
| Samburu County         | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | •30                          | 30%  | 0%                                  | 5                                    | 74  | 100%  | 55%  | 96%  | 0 42   |  |
| Siaya County           | Lake endemic                | <b>*</b> 119  | <b>*</b> 104  | <b>*</b> 78          | <b>4</b> 4                   | <b>*</b> 81%   | <b>↓</b> 32%                        | <b>↓</b> 155                         | ♦95   | 0%  | <b>€</b> 52%   | 99%  | 46.66  |  |
| Taita Taveta County    | Coast endemic               | <b>1</b> 07   | <b>*</b> 91   | €94                  | <b>*</b> 36                  | 100%   | 0%                                  | 1                                    | 58  | 0%  | <b>1</b> 90%   | <b>*</b> 96%   | 0.88   |  |
| Tana River County      | Seasonal, arid, & semi-arid | €68   | ♦69   | ●32                  | •33                          | <b>₽</b> 32%   | <b>₽</b> 3%                         | 4                                    | €128  | 100%  | 43%  | <b>↑</b> 84%   | 0 7 1  |  |
| Tharaka Nithi County   | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | €42                          | <b>₽</b> 35%   | <b>€</b> 2%                         | 1                                    | <b>●</b> 31   | 100%  | 94%  | <b>€</b> 95%   | 0 7  |  |
| Trans Nzoia County     | Highland epidemic prone     | <b>*</b> 85   | <b>†</b> 92   | 0                    | 24                           | <b>₽</b> 21%   | <b>*</b> 4%                         | ₹22                                  | <b>•</b> 75   | 100%  | 70%  | <b>1</b> 94%   | 1.76   |  |
| Turkana County         | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | <b>\$</b> 35                 | <b>\$</b> 59%  | <b>1</b> 7%                         | ≢51                                  | <b>\$</b> 89  | 100%  | <b>*</b> 94%   | 89%  | 2.59   |  |
| Uasin Gishu County     | Highland epidemic prone     | <b>₽</b> 72   | ₹71   | 0                    | 38                           | 100%   | 2%                                  | 7                                    | 78  | 100%  | €89%   | <b>1</b> 99%   | 0  |  |
| Vihiga County          | Lake endemic                | ♦107  | <b>1</b> 31   | <b>₽</b> 79          |                              | 84%  | <b>€</b> 53%                        | <b>\$</b> 146                        |   | 0%  | ●95%   | 100%   | 0.86   |  |
| Wajir County           | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | €28                          | <b>*</b> 85%   | 0%                                  | 0                                    | <b>€</b> 657  | 100%  | <b>●</b> 87%   | <b>↑</b> 86%   | 0  |  |
| West Pokot County      | Seasonal, arid, & semi-arid | 0   | 0   | 0                    | 21                           | <b>*</b> 53%   | 4%                                  | ♦30                                  | ♦95   | 100%  | 77%  | 91%  | 0 –  |  |
| Source:                | Policy                      | DHIS  | DHIS  | DHIS                 | DHIS                         | DHIS   | DHIS                                | DHIS                                 | DHIS  | NMCP  |  | DHIS   | Activity Report /  |  |