

Country indicators

% HHS that have attained UC of LLINs in targeted areas	% PW who slept under an LLIN among HHS with at least 1 net	% of pregnant women who received 3+ doses of IPTp for malaria during their last pregnancy in the last 2 yrs	% HHS having no stock-out of ACTs	ACTs distributed against projected Forecasting & Quantification targets	% patients treated according to national guidelines	Malaria prevalence in children <5 yrs
51%	85%	83%	38%	71%	81%	100%
DHS/KMIS	DHS/KMIS / DHS/KMIS	KMIS	QOC survey / QOC Survey	KEMSA	QOC Survey	KMIS

Scorecard

Region	Malaria Prevention			Malaria in Pregnancy		Case Management				EPR	Surveillance Monitoring & Evaluation and Operational Research			
	Transmission Zone	% PW receiving an LLIN at ANC clinic in targeted areas	% children <1 yr receiving an LLIN at immunization clinic in targeted areas	% PW who received IPT 2 in ANC malaria endemic areas	% PW who had 4 ANC visits	% suspected malaria cases receiving a parasitological test	Annual Blood Examination Rate	Malaria cases per 1000 population	Outpatient cases treated w/ AL as a proportion of confirmed malaria cases	% epidemic prone and seasonal transmission sub counties with updated annual EPR plans	Reporting rates of MOH 505 (weekly surveillance)	Reporting rate for malaria commodity form	Annual entomological inoculation rate / Confirmed Malaria Cases per 1,000 Population per Year	
Kenya		62	63	68	37	77%	11%	26	87	100%	76%	93%	4.29	
Baringo	Seasonal, arid, & semi-arid	0	0	0	27	46%	3%	11	123	100%	52%	89%	0.83	
Bomet	Highland epidemic prone	69	84	0	29	38%	1%	1	159	100%	79%	85%	0	
Bungoma	Lake endemic	74	72	78	43	83%	41%	85	85	100%	91%	100%	0	
Busia	Lake endemic	81	81	104	39	91%	86%	152	84	0%	95%	100%	33.97	
Elgeyo-Marakwet	Seasonal, arid, & semi-arid	0	0	0	23	88%	0%	6	57	100%	88%	86%	0	
Embu	Seasonal, arid, & semi-arid	0	0	0	37	98%	0%	0	62	100%	75%	99%	0	
Garissa	Seasonal, arid, & semi-arid	0	0	0	67	63%	1%	2	37	100%	77%	93%	0	
Homa Bay	Lake endemic	62	59	64	36	57%	10%	32	23	0%	81%	100%	7.32	
Isiolo	Seasonal, arid, & semi-arid	0	0	0	35	57%	1%	4	84	100%	62%	76%	0	
Kajiado	Seasonal, arid, & semi-arid	0	0	0	37	37%	0%	1	55	100%	85%	93%	0	
Kakamega	Lake endemic	106	102	69	41	83%	77%	136	99	100%	99%	100%	6.33	
Kericho	Highland epidemic prone	64	57	0	27	77%	5%	8	79	100%	96%	96%	0.99	
Kiambu	Low risk	0	0	0	41	63%	0%	0	27	0%	80%	88%	0	
Kilifi	Coast endemic	76	68	52	35	38%	7%	13	81	0%	51%	96%	1.81	
Kirinyaga	Low risk	0	0	0	42	73%	1%	0	59	0%	72%	98%	3.01	
Kisii	Highland epidemic prone	63	68	0	30	74%	7%	10	76	100%	95%	96%	9.94	
Kisumu	Lake endemic	95	94	63	45	83%	31%	87	91	0%	52%	92%	13.86	
Kitui	Seasonal, arid, & semi-arid	0	0	0	33	30%	1%	0	50	100%	92%	91%	0	
Kwale	Coast endemic	78	79	70	37	27%	4%	37	120	0%	31%	96%	0.8	
Laikipia	Low risk	0	0	0	38	58%	0%	1	17	0%	71%	99%	0	
Lamu	Coast endemic	96	82	90	46	37%	0%	0	60	0%	15%	97%	0	
Machakos County	Low risk	0	0	0	45	100%	0%	0	100	0%	100%	98%	0	
Makueni	Low risk	0	0	0	51	26%	4%	0	17	0%	100%	100%	0	
Mandera	Seasonal, arid, & semi-arid	0	0	0	30	0%	0%	0	76	100%	0%	67%	0	
Marsabit	Seasonal, arid, & semi-arid	0	0	0	33	76%	0%	2	34	100%	27%	87%	0	
Meru	Seasonal, arid, & semi-arid	0	0	0	22	0%	0%	0	65	100%	37%	31%	0	
Migori	Lake endemic	98	92	63	37	79%	38%	80	86	0%	92%	100%	14.25	
Mombasa	Coast endemic	59	61	76	37	54%	7%	4	23	0%	77%	94%	0	
Muranga	Low risk	0	0	0	39	56%	0%	0	4	0%	62%	96%	0	
Nairobi	Low risk	0	0	0	45	66%	1%	2	28	0%	51%	87%	0	
Nakuru	Low risk	0	0	0	36	61%	0%	2	34	0%	75%	85%	0	
Nandi	Highland epidemic prone	113	150	0	29	79%	6%	16	93	100%	70%	91%	3.35	
Narok	Highland epidemic prone	85	98	0	27	31%	5%	4	132	100%	77%	78%	0	
Nyamira	Highland epidemic prone	76	71	0	51	43%	2%	10	60	100%	74%	97%	1.04	
Nyandarua	Low risk	0	0	0	35	100%	0%	0	19	0%	69%	100%	0	
Nyeri	Low risk	0	0	0	42	27%	0%	0	52	0%	77%	100%	0	
Samburu	Seasonal, arid, & semi-arid	0	0	0	30	30%	0%	5	74	100%	55%	96%	0	
Siaya	Lake endemic	119	104	78	44	81%	32%	155	95	0%	52%	99%	46.66	
Taita Taveta	Coast endemic	107	91	94	36	100%	0%	1	58	0%	90%	96%	0.88	
Tana River	Seasonal, arid, & semi-arid	68	69	32	33	32%	3%	4	128	100%	43%	84%	0	
Tharaka Nithi	Seasonal, arid, & semi-arid	0	0	0	42	35%	2%	1	31	100%	94%	95%	0	
Trans-Nzoia	Highland epidemic prone	85	92	0	24	21%	4%	22	75	100%	70%	94%	1.76	
Turkana	Seasonal, arid, & semi-arid	0	0	0	35	59%	17%	51	89	100%	94%	89%	2.59	
Uasin Gishu	Highland epidemic prone	72	71	0	38	100%	2%	7	78	100%	89%	99%	0	
Vihiga	Lake endemic	107	131	79	39	84%	53%	146	71	0%	95%	100%	0.86	
Wajir	Seasonal, arid, & semi-arid	0	0	0	28	85%	0%	0	657	100%	87%	86%	0	
West Pokot	Seasonal, arid, & semi-arid	0	0	0	21	53%	4%	30	95	100%	77%	91%	0	
Source:	Policy	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	DHIS	NMCP	DHIS	DHIS	Activity Report / DHIS	